



**GASTROINTESTINAL (GI) Requisition to PET Centre**  
**TO BE COMPLETED BY THE REFERRING PHYSICIAN**

**Indications:** (choose only one)

**Patient Name:** \_\_\_\_\_

- COLORECTAL CANCER (STAGING/RE-STAGING) – PET for patients with apparent limited metastatic disease (e.g., organ-restricted liver or lung metastases; or limited local recurrence) who are being considered for radical intent therapy.**

**Note:** As chemotherapy may affect the sensitivity of the PET scan, it is strongly recommended to schedule PET at least 6 weeks after last chemotherapy, if possible.

**The patient must have:**

- Histologic confirmation of colorectal cancer, **AND**  
 Presumptive pre-PET apparent limited metastatic disease or local recurrence, **AND**  
 Patient has **no** significant comorbidities that would preclude radical intent therapy, if clinically indicated.

Attach the relevant diagnostic imaging reports; and provide images to the PET Centre.

**Other information regarding eligibility:** \_\_\_\_\_

- COLORECTAL CANCER (RECURRENCE) – PET where recurrent disease is suspected on the basis of elevated and/or rising carcinoembryonic antigen (CEA) level(s) after surgical resection with negative or equivocal imaging work-up.**

**The patient must have:**

- Received primary therapy, **AND**  
 Recent imaging (CT or MRI) that is **negative or equivocal**, **AND**  
 Elevated Biomarker: Biomarker : \_\_\_\_\_ Value 1 : \_\_\_\_\_ Value 2 : \_\_\_\_\_

Attach the most recent biomarker results & the relevant diagnostic imaging reports; and provide images to the PET Centre.

**Other information regarding eligibility:** \_\_\_\_\_

- ANAL CANAL CANCER (STAGING/RE-STAGING) - PET for the initial staging of patients with clinical stage II-IV squamous cell carcinoma of the anal canal or when conventional imaging is equivocal for a specific stage; or for re-staging of patients with limited recurrence, after primary treatment, being considered for definitive salvage therapy.**

**Purpose of PET scan (choose 1):**

- Initial staging of patients with clinical stage II-IV SCC of the anal canal  
 Initial staging to clarify equivocal conventional imaging of patients with SCC of the anal canal, specify location(s) of interest for PET:  
 Ano-rectum     Lymph Nodes     Elsewhere (specify): \_\_\_\_\_  
 Re-staging (limited recurrence) when further ablative therapy is being considered

Attach relevant diagnostic imaging reports (CT, US, MR, Other) & provide images to the PET Centre); consult note or referral letter; and the pathology report.

**Physician Signature:** \_\_\_\_\_ **Date:** \_\_\_\_\_